APS/123-QED

arXiv:1209.5326v1 [cond-mat.stat-mech] 24 Sep 2012

Granular gas of viscoelastic particles in a heated and force-free state

Awadhesh Kumar Dubey,[1] Anna Bodrova,[2] Sanjay Puri,[1] and Nikolai Brilliantov[3]

[1] *School of Physical Sciences, Jawaharlal Nehru University, New Delhi, 110067, India.*
[2] *Faculty of Physics, M.V.Lomonosov Moscow State University, Moscow, 119991, Russia*
[3] *Department of Mathematics, University of Leicester, Leicester LE1 7RH, United Kingdom*

(Dated: September 25, 2012)

We present large-scale Molecular Dynamics simulation results for granular gases of particles with the velocity-dependent restitution coefficient, as it follows from the simplest first-principle model of viscoelastic spheres. We analyze both cases of force-free gases and homogeneously heated gases. We develop a theory for the velocity distribution for heated gases. We explored evolution of temperature and the Sonine coefficient, which characterize the form of the velocity distribution function. We observed that for not large dissipation the simulation results are in excellent agreement with both, the existing theory for gases in a homogeneous cooling state and with the novel theory for heated gases. At the same time a noticeable discrepancy between theory and MD results have been detected for large dissipations. We analyze the adequacy of the simplified model of a constant effective restitution coefficient for uniformly heated gases of viscoelastic particles. We conclude that while this model may be used to describe temperature of a gas, it fails to describe quantitatively both temperature and the velocity distribution function.

PACS numbers: 81.05.Rm, 05.20.Dd, 05.40.2a

I. INTRODUCTION

Granular materials are abundant in nature and examples range from sands and powders on Earth [1–5] to more dilute systems, termed as granular gases [6–9] in space; astrophysical objects, like planetary rings, may be mentioned in this respect [6, 10]. These systems are of a great scientific and industrial importance. Depending on the average kinetic energy of individual grains, granular materials can be in a solid, liquid or gaseous state. In this paper, we address a low-density granular gas – a system comprised of macroscopically large number of macroscopic grains that move ballistically between inelastic pair-wise collisions. The inelastic nature of inter-particle collisions is the main feature which distinguishes a granular gas from a molecular gas and gives rise to many astonishing properties of dissipative gases. If no external forces act on the system it evolves freely and permanently cools down. During the first stage of a granular gas evolution its density remains uniform and no macroscopic fluxes present. This state, which is called a *homogeneous cooling state* (HCS) [8, 11], is however, unstable with respect to fluctuations and in later times clusters and vertexes develop. The system then continue to loose its energy in the *inhomogeneous cooling state* (ICS) [12–19]. If energy is injected into a granular gas to compensate the loss of kinetic energy, the system settles into a nonequilibrium steady-state [20, 21].

The dissipation of energy in pair-wise collisions is quantified by the restitution coefficient ε,

$$\varepsilon = \left| \frac{(\mathbf{v}_{12}' \cdot \mathbf{e})}{(\mathbf{v}_{12} \cdot \mathbf{e})} \right| , \tag{1}$$

where $\mathbf{v}_{12}' = \mathbf{v}_2' - \mathbf{v}_1'$ and $\mathbf{v}_{12} = \mathbf{v}_2 - \mathbf{v}_1$ are the relative velocities of two particles after and before a collision, and $\mathbf{e}$ is a unit vector connecting their centers at the collision instant. The post-collision velocities are then related to the pre-collision velocities $\mathbf{v}_1$ and $\mathbf{v}_2$ as follows, e.g. [8]:

$$\mathbf{v}_{1/2}' = \mathbf{v}_{1/2} \mp \frac{1+\varepsilon}{2}(\mathbf{v}_{12} \cdot \mathbf{e})\mathbf{e} \, , \tag{2}$$

To date, most studies of the HCS and ICS have focused on the case of a constant restitution coefficient [11–19]. This assumption, however, contradicts experimental observations [22–24], along with basic mechanical laws [25, 26], which show that ε does depend on the impact velocity [24, 26–29]. This dependence may be obtained solving equation of motion for colliding particles with the explicit account for the dissipative forces acting between the grains. The simplest first-principle model of inelastic collisions assumes viscoelastic properties of particles material, which result in viscoelastic inter-particle force [27] and finally in the restitution coefficient [26, 29]:

$$\varepsilon = 1 - C_1 A \kappa^{2/5} \left| \mathbf{v}_{12} \cdot \mathbf{e} \right|^{1/5} + C_2 A^2 \kappa^{4/5} \left| \mathbf{v}_{12} \cdot \mathbf{e} \right|^{2/5} \; + \ldots \, . \tag{3}$$

Here, the numerical coefficients are $C_1 \simeq 1.15$, $C_2 \simeq 0.798$. The elastic constant

$$\kappa = \left(\frac{3}{2} \right)^{3/2} \frac{Y \sqrt{\sigma}}{m(1-\nu^2)} \tag{4}$$

is a function of the Young's modulus Y and Poisson ratio ν and mass m; the constant A quantifies the viscous properties of the particles' material. While the last equation is valid for not very large inelasticities, in the recent work of Schwager and Poschel [30] an expression for ε, which is valid for highly inelasticity and accounts for the delayed recovery of the shape of colliding particles has been derived:

$$\varepsilon = 1 + \sum_{k=1}^{N_k} h_k \delta^{k/2} \left[2u\left(t\right) \right]^{k/20} \left| (\mathbf{c}_{12} \cdot \mathbf{e}) \right|^{k/10} \, . \tag{5}$$

Here, $h_1 = 0$, $h_2 = -C_1$, $h_3 = 0$, $h_4 = C_2$, and the other numerical coefficients up to $N_k = 20$ are given in Ref. [30]. In the latter equation $u(t) = T(t)/T_0$ ($T_0 = T(0)$) is the reduced temperature, with the usual definition of temperature

$$\frac{3}{2}nT(t) = \int d\mathbf{v}\ \frac{m\mathbf{v}^2}{2} f(\mathbf{v},t)\,. \tag{6}$$

Here n is the number density of the gas, $\mathbf{v}$ is a particle velocity, $f(\mathbf{v},t)$ – the time-dependent velocity distribution function and $\mathbf{c}_{12}$ is the dimensionless relative velocity, defined as $\mathbf{c}_{12} = \mathbf{v}_{12}/v_T$, where $v_T(t) = \sqrt{2T(t)/m}$ is the thermal velocity. The dissipative constant δ reads,

$$\delta = A\kappa^{2/5}\left(\frac{T_0}{m}\right)^{1/10}, \tag{7}$$

so that the first few terms in Eq. (5) are identical to those in Eq. (3). Note that $\delta = 0$ corresponds to an elastic system, $\varepsilon = 1$ and the dissipation in the system increases with increasing δ. Furthermore, $\varepsilon \to 1$ when the reduced temperature $u(t)$ decreases, that is, the effective restitution coefficient increases as a system cools down. The dependence of ε on the dissipation parameter δ is shown in Fig. 1 for $u(t) = 1$ and the average value of the reduced normal velocity $\langle|\mathbf{c}_{12}\cdot\mathbf{e}|\rangle = 4/\sqrt{2\pi}$. As it may be seen from the figure, the restitution coefficient, given by Eq. (5) demonstrates a good convergence for $\delta \lesssim 0.5$. In what follows we use the number of terms in the expansion (5) equal to $N_k = 20$, which guarantees an accurate description of ε up for $\delta \lesssim 0.5$, that is, for rather large dissipation.



FIG. 1: Normal restitution coefficient ε, as it follows from from Eq. (5), for different values of N_k. Here $u = 1$ and $\langle|\mathbf{c}_{12}\cdot\mathbf{n}|\rangle = 4/\sqrt{2\pi}$. The latter value is the average dimensionless relative normal impact velocity with the averaging performed over a Maxwellian distribution.

The velocity distribution function in granular gases usually deviates from the Maxwellian distribution [18, 19, 31–33] and may be represented in terms of a *Sonine-polynomial* expansion [8]:

$$f(\mathbf{v},t) = \frac{n}{v_T^3}\tilde{f}(\mathbf{c},t),$$
$$\tilde{f}(\mathbf{c},t) = \pi^{-3/2}\exp\left(-c^2\right)\left[1 + \sum_{p=1}^{\infty} a_p(t)S_p(c^2)\right], \tag{8}$$

where the Sonine polynomials read for $d = 3$:

$$S_p(c^2) = \sum_{k=0}^{p} \frac{(-1)^k(1/2+p)!}{(1/2+k)!(p-k)!k!}c^{2k}\,. \tag{9}$$

Hence evolution of $f(\mathbf{v},t)$ is completely determined by the time dependence of the Sonine coefficients $a_p(t)$. The first Sonine coefficient is trivial, $a_1 = 0$, as it follows from the definition of temperature, e.g. [8] while the elastic Maxwellian limit corresponds to $a_p = 0$ for all $p > 1$.

It has been shown recently that for granular gases with a constant ε the expansion (8) converges for small and moderate dissipation, up to $\varepsilon \gtrsim 0.6$, but breaks down otherwise [34]. If the dissipation is not large, one can consider only the first two non-trivial coefficients a_2 and a_3, which have been studied both analytically [31, 33, 34] and numerically [15, 18, 19, 34–36].

The theory developed for the case of constant ε predicts a rapid relaxation of a_p to steady state values, which depend on ε [33, 34]. The solution of the Boltzmann equation by the Direct Simulation Monte Carlo (DSMC) method demonstrated a good agreement between theoretical and numerical results [34]. The analysis of the granular gas dynamics by the DSMC has however a serious drawback – it does not take into account correlations between particles velocities which arise during system's evolution. It is not known *a priori* whether these correlations have an impact on the studied gas properties. Indeed, Molecular dynamics (MD) simulations which are free from the above shortcoming show at certain densities noticeable deviations of the numerically obtained coefficients $a_p(t)$ from the corresponding theoretical values [18, 19]. This discrepancy arises due to formation of spatial structures in a gas, which spontaneously appear when the gas evolves in a HCS.

Evolution of the velocity distribution function and respectively the Sonine coefficients is qualitatively different in a gas of viscoelastic particles. In contrast to the case of a constant ε, where rapid relaxation is observed, the evolution of the Sonine coefficients a_2 and a_3 demonstrates a complicated non-monotonous time dependence [37, 38]. Typically, the magnitudes of the Sonine coefficients first increase with time, reach maximal values, and finally tend to zero, so that asymptotically the Maxwellian distribution is recovered [37, 38]. This simply follows from the fact that the collisions become more and more elastic as the average particles velocity decreases [cf. Eq. (5)]. A theory describing evolution of $a_2(t)$ and $a_3(t)$ in a force-free granular gas of viscoelastic particles has been developed in [38], however a corresponding theory for heated gases is still lacking. Moreover, the behavior of the velocity distribution function in a gas of viscoelastic particles has never been studied by means of MD neither for force-free nor for driven systems. Hence, it is still not known how accurate is the theory for the velocity distribution function in this case and for which dissipation δ the Sonine series keeps its convergence. Another interesting question is the adequacy of the simplified model of a constant restitution coefficient for uniformly heated gases of viscoelastic particles: While it is known that model of constant ε fails to describe a complicated evolution of a gas in a homogeneous cooling state [40], it is still not known how good this model for driven systems.

In the present study we perform thorough large-scale MD simulation studies of the velocity distribution function in a gas of viscoelastic particles, both in a homogeneous cooling state and under a uniform heating; we also develop a theory for $f(\mathbf{v},t)$ for driven gases. We observe that our MD results for $a_2(t)$ and $a_3(t)$ are in excellent agreement with the analytical predictions of Ref. [38] for the HCS, as well as with the results of the novel theory for driven systems, reported here. We present also simulation results for the Sonine coefficients a_4 and a_5, which may be hardly obtained from the kinetic theory. The analysis of the series of these coefficients, $a_2,\ a_3,\ a_4,\ a_5$, allows to asses the convergence of the Sonine-polynomial expansion. The problem of an adequacy of the model of the effective constant restitution coefficient for driven gases of viscoelastic particles is also addressed.

This paper is organized as follows. In the next Sec. II we develop a theory for evolution of the velocity distribution function in a heated gas of viscoelastic particles, that is, we obtain the time dependence of the Sonine coefficients; the previous results for $a_2(t)$ and $a_3(t)$ for a force-free gas are directly reproduced in a limiting case of vanishing heating. In Sec. III we report our MD results and compare simulation data with the theoretical predictions. Finally, in Sec. IV we summarize our findings.

II. KINETIC THEORY OF UNIFORMLY HEATED GRANULAR GAS OF VISCOELASTIC PARTICLES

We address a granular gas of identical viscoelastic particles in a white-noise thermostat, that is, the gas under a uniform heating. The equation of motion for i-th particle then reads:

$$m\frac{d\mathbf{v}_i}{dt} = \mathbf{F}_i^{\text{coll}} + \mathbf{F}_i^{\text{ext}}\,. \tag{10}$$

Here the force acting between colliding i-th and j-th particles is of the form [27]

$$\mathbf{F}_i^{\text{coll}} = \sum_j \left(\rho\zeta_{ij}^{3/2} + \frac{3}{2}A\rho\zeta_{ij}^{1/2}\frac{d\zeta_{ij}}{dt}\right)\Theta(\zeta_{ij})\mathbf{e}_{ij}, \tag{11}$$

where $\rho = \frac{1}{2}m\kappa\,(2/3)^{5/2}$ and $\zeta_{ij} = r_{ij} - \sigma$ is the deformation of particles in a pairwise collision. It is expressed in terms of the inter-particle distance, $r_{ij} = |\mathbf{r}_i - \mathbf{r}_j|$ and particle's diameter σ. The unit vector $\mathbf{e}_{ij} = \mathbf{r}_{ij}/r_{ij}$ gives the direction of the inter-particle force and the Heaviside step-function $\Theta(x)$ indicates that the force acts only at a collision, when particles' centers approach closer than the diameter σ. In Eq. (11) the summation over all grains is performed. For the external force $\mathbf{F}^{\text{ext}}$ we use the model of a Gaussian white noise:

$$\langle\mathbf{F}_i^{\text{ext}}(t)\rangle = 0, \quad \langle F_{i,\alpha}^{\text{ext}}(t)F_{j,\beta}^{\text{ext}}(t')\rangle = m^2\xi_0^2\delta_{ij}\delta_{\alpha\beta}\delta(t-t')\,, \tag{12}$$

where $\alpha,\beta = x,y,z$, and ξ_0^2 characterizes the magnitude of the stochastic force.

We use event-driven simulations [39], that is, we do not explicitly solve an equation of motion for colliding particles with the forces $\mathbf{F}^{\text{coll}}$, but directly use the after-collision velocities, expressed in terms of the pre-collision ones and the coefficient of restitution ε [8, 39]. If we further apply the hypothesis of molecular chaos, evolution of the velocity distribution function may be described using the Boltzmann-Enskog (BE) equation, supplemented by the diffusive Fokker-Planck term, which corresponds to a uniform heating [21]:

$$\frac{\partial f(\mathbf{v},t)}{\partial t} = g_2(\sigma)I(f,f) + \frac{\xi_0^2}{2}\frac{\partial^2}{\partial\mathbf{v}^2}f(\mathbf{v},t). \tag{13}$$

Here $I(f,f)$ is the collision integral and $g_2(\sigma)$ is contact value of the pair correlation function that takes into account the excluded volume effect [8]. For the scaling distribution function $\tilde{f}(\mathbf{c},t)$ [cf. Eq. (8)], the above equation reads

$$\frac{\partial\tilde{f}}{\partial t} - \frac{1}{v_T}\frac{dv_T}{dt}\left(3\tilde{f} + c\frac{\partial\tilde{f}}{\partial c}\right) = g_2\sigma^2 n v_T\tilde{I} + \frac{1}{v_T^2}\frac{\xi_0^2}{2}\frac{\partial^2}{\partial\mathbf{c}^2}\tilde{f}. \tag{14}$$

Here $\tilde{I}\left(\tilde{f},\tilde{f}\right) = \sigma^{-2}n^{-2}v_T^2 I\left(f,f\right)$ is the dimensionless collision integral, defined as

$$\tilde{I}(\tilde{f},\tilde{f}) \;=\; \int d\mathbf{c}_2 \int d\mathbf{e}\,\Theta(-\mathbf{c}_{12}\cdot\mathbf{e}\,)\,|-\mathbf{c}_{12}\cdot\mathbf{e}\,| \qquad (15)$$
$$\times\;\left[\frac{1}{\varepsilon^2}\tilde{f}(\mathbf{c}_1{''},t)\tilde{f}(\mathbf{c}_2{''},t)-\tilde{f}(\mathbf{c}_1,t)\tilde{f}(\mathbf{c}_2,t)\right]\,.$$

Here $\mathbf{c}_1''$ and $\mathbf{c}_2''$ are the (reduced) pre-collision velocities in the so-called inverse collision, which results in the (reduced) post-collision velocities $\mathbf{c}_1$ and $\mathbf{c}_2$ [8].

Now we substitute the Sonine-polynomial expansion for $\tilde{f}(\mathbf{c},t)$ [Eq. (8)] into Eq. (14) and neglect all terms with $p>3$. Multiplying the resulting equation with c_1^2, c_1^4 and c_1^6, and integrating over $\mathbf{c}_1$, we obtain

$$\begin{aligned}
\frac{\partial\langle c^2\rangle}{\partial t}+\frac{1}{T}\frac{dT}{dt}\langle c^2\rangle &= -\sqrt{\frac{2T}{m}}g_2(\sigma)\sigma^2 n\mu_2+3\frac{m\xi_0^2}{2T}, \qquad (16)\\
\frac{\partial\langle c^4\rangle}{\partial t}+\frac{2}{T}\frac{dT}{dt}\langle c^4\rangle &= -\sqrt{\frac{2T}{m}}g_2(\sigma)\sigma^2 n\mu_4+10\langle c^2\rangle\frac{m\xi_0^2}{2T},\\
\frac{\partial\langle c^6\rangle}{\partial t}+\frac{3}{T}\frac{dT}{dt}\langle c^6\rangle &= -\sqrt{\frac{2T}{m}}g_2(\sigma)\sigma^2 n\mu_6+24\langle c^4\rangle\frac{m\xi_0^2}{2T}.
\end{aligned}$$

Here,

$$\mu_p=-\int d\mathbf{c}c^p\tilde{I}(\tilde{f},\tilde{f}) \qquad (17)$$

is the p-th moment of the dimensionless collision integral. The moments μ_p can be calculated analytically up to $\mathcal{O}(\delta^{10})$ using a formula manipulation program as explained in detail in Ref. [8]. They can be represented in the following form:

$$\begin{aligned}
\mu_p \;=\; &\sum_{k=0}^{20}\Big(M_{p,\,0}^{(k)}+M_{p,\,2}^{(k)}a_2+M_{p,\,3}^{(k)}a_3+ \qquad (18)\\
+\;&M_{p,\,22}^{(k)}a_2^2+M_{p,\,33}^{(k)}a_3^2+M_{p,\,23}^{(k)}a_2a_3\Big)\,\delta^{k/20}\,(2u)^{k/20}\,.
\end{aligned}$$

The numerical values of the coefficients $M_{p,\,i}^{(k)}$ for μ_p with $p=2,\,4,\,6$ are given in Appendix.

The Sonine coefficients are related to the moments of the reduced velocity as follows:

$$\begin{aligned}
\left\langle c^2\right\rangle &= \frac{3}{2}, \qquad (19)\\
\left\langle c^4\right\rangle &= \frac{15}{4}\left(1+a_2\right),\\
\left\langle c^6\right\rangle &= \frac{105}{8}\left(1+3a_2-a_3\right).
\end{aligned}$$

Hence using Eqs. (16), one can obtain the following system of coupled equations:

$$\begin{aligned}
\frac{du}{d\tau} &= -\frac{\sqrt{2}\mu_2}{6\sqrt{\pi}}u^{\frac{3}{2}}+\frac{K}{4}, \qquad (20)\\
\frac{da_2}{d\tau} &= \frac{\sqrt{2}\sqrt{u}}{3\sqrt{\pi}}\mu_2\left(1+a_2\right)-\frac{\sqrt{2}}{15\sqrt{\pi}}\mu_4\sqrt{u}-\frac{Ka_2}{2u},\\
\frac{da_3}{d\tau} &= \frac{\sqrt{u}}{\sqrt{2\pi}}\mu_2(1-a_2+a_3)-\\
&- \frac{\sqrt{2}}{5\sqrt{\pi}}\mu_4\sqrt{u}+\frac{2\sqrt{2u}}{105\sqrt{\pi}}\mu_6-\frac{3a_3K}{4u}\,,
\end{aligned}$$

with the constant K,

$$K=\frac{m^{3/2}\xi_0^2}{\sqrt{\pi}g_2(\sigma)\sigma^2 nT_0^{3/2}}. \qquad (21)$$

In the above equation we use the dimensionless time $\tau=t/\tau_c(0)$, expressed in terms of the initial mean collision time $\tau_c(0)$, where

$$\tau_c^{-1}(t)=4\sqrt{\pi}g_2(\sigma)\sigma^2 n\sqrt{\frac{T(t)}{m}}. \qquad (22)$$

Eqs. (20) describe evolution of temperature and the velocity distribution function for a heated gas of viscoelastic particles. For $K=0$, that is for $\xi_0^2=0$, Eqs. (20) reduce to corresponding equations of Ref. [38], which describe evolution of a_2 and a_3 in a force-free gas. Note that Eqs. (20) also describe evolution of Sonine coefficients for a granular gas with a constant restitution coefficient. In this case the respective values of μ_p for a constant ε are to be used Ref. [34].

When a heated gas settles in a steady state, the granular temperature and the velocity distribution function become time-independent. Then Eqs. (20) yield the algebraic equations for the steady-state temperature and moments μ_p, $p=2$, 4, 6:

$$\begin{aligned}
T &= \left[\frac{3}{2\sqrt{2}}\frac{\xi_0^2 m^{3/2}}{\mu_2 g_2(\sigma)\sigma^2 n}\right]^{2/3}, \qquad (23)\\
\mu_4 &= 5\mu_2, \qquad (24)\\
\mu_6 &= \frac{105}{4}\mu_2(1+a_2). \qquad (25)
\end{aligned}$$

Solving this system in the linear approximation with respect to the dissipation parameter δ, one obtains the steady state Sonine coefficients

$$a_2^{\text{s.s.}}=-A_2\delta, \qquad A_2=2^{1/5}\frac{157}{500}\Gamma\left(\frac{21}{10}\right)C_1\simeq 0.435, \qquad (26)$$

$$a_3^{\text{s.s.}}=-A_3\delta, \qquad A_3=2^{1/5}\frac{28}{500}\Gamma\left(\frac{21}{10}\right)C_1\simeq 0.078. \qquad (27)$$

The above expressions for a_2 and a_3, together with Eq. (19) for μ_2 yield the corresponding analytical expression for the steady state temperature $T_{\text{s.s.}}$.

III. MOLECULAR DYNAMICS SIMULATIONS

A. Force-free Gas

We perform MD simulations [41], using an event-driven algorithm [39] to investigate a system of inelastic granular particles with the restitution coefficient (5), as it follows for viscoelastic particles. The grains are modeled as identical hard spheres of mass $m = 1$ and diameter $\sigma = 1$. In event-driven simulations particles move freely between pairwise collisions, where the velocities of the particles are updated according to Eq. (2). Our system consists of $N = 2.048 \times 10^6$ particles placed in a three-dimensional cubic box with the edge $L = 418$, which corresponds to the number density of $n = 0.028$. Periodic boundary conditions are applied in all three directions.

The system was initialized by placing randomly particles in the box. For initial particles' velocities randomly directed vectors of the same length have been assigned. The initial total momentum $\sum_i m\mathbf{v}_i(0)$ was in this case zero with a high accuracy. We start with $\varepsilon = 1$, which corresponds to elastic collisions, and allow the system to relax to a Maxwellian velocity distribution, which used to happen in a few collisions per particle. Hence, a homogeneous system with a Maxwellian distribution was an initial condition for each MD run; we then evolve the system with a velocity-dependent restitution coefficient (5). In simulations we used the initial granular temperature $T(0) = 400/3$ with the corresponding initial mean collision time $\tau_c(0) = 0.436$. We have confirmed that the system remains approximately homogeneous during our simulations. All statistical quantities presented here have been obtained as averages over 25 independent runs.

In Fig. 2, we present simulation results for the dependence of the reduced granular temperature $u(\tau)$ on the dimensionless time τ. In a HCS temperature decays according to the power-law $u(\tau) \sim \tau^{-5/3}$ [8, 37, 38]. The agreement between MD results (denoted by symbols) and theory (denoted by lines) is very good for both small and moderate values of the dissipative parameter δ; as expected, the larger δ, the faster the gas cools. In Fig. 2, we also show the results for the heated gas, which are discussed in the following subsection. Next, we analyze evolution of the velocity distribution, characterized by the time-dependent Sonine coefficients [cf. Eq. (8)]. In Fig. 3, we show evolution of a_2 and a_3 for a force-free gas with $\delta = 0.15$. In order to calculate a_2 and a_3 numerically, we compute the moments of the velocity distribution function $\langle c^{2k} \rangle$ and use Eqs. (19). At initial time, the Sonine coefficients equal to zero: $a_2(0) = a_3(0) = 0$, which corresponds to a Maxwellian distribution. As it follows from Fig. 3, the absolute values of a_2 and a_3 initially increase, reach a maximum, and eventually decay to zero. As the thermal velocity decreases during the gas cooling, the behavior of the system tends to that of a gas of elastic particles, that is, the velocity distribution tends to a Maxwellian. The maximum values of $|a_2|$ and $|a_3|$ increase with δ, saturate at $\delta \simeq 0.15$, and do not increase further; the location of these maxima shifts to later times with increasing δ [38]. To investigate the convergence of the Sonine-polynomial expansion (8) we analyze the high-order Sonine coefficients, a_4 and a_5. In Fig. 4 the simulation data for the time dependence of a_2, a_3, a_4, a_5 along with the analytical predictions for a_2 and a_3 are shown for $\delta = 0.1$. Note, that a_4 and a_5 evolve in a similar fashion as a_2 and a_3, that is, their magnitudes first increase, reach maximum and relax back to zero. Certainly, this is not surprising, since it is a simple manifestation of the fact that the system starts from the Maxwellian velocity distribution due to initial conditions and returns to it later, when the collisions become elastic. Again we see an excellent agreement between the simulation results and the theory for a_2 and a_3. Furthermore, a_4 is an order of magnitude smaller than a_2, while a_5 is $\sim 10^2$ times smaller than a_2. This confirms the convergence of the Sonine-polynomial expansion for the velocity distribution, at least for these values of δ.



FIG. 2: Evolution of the reduced granular temperature $u(\tau) = T(\tau)/T(0)$ for the force-free and heated granular gases. The symbols denote simulation results, the lines – theoretical predictions [the numerical solution of Eqs. (20)]. Temperature of the force-free gas decreases asymptotically, $\tau \gg 1$, as $u \sim \tau^{-5/3}$, while that of the heated gas first decreases and then saturates at a constant, steady state value $u_{\text{s.s.}}$. Gray lines correspond to evolution of granular gas with the effective constant restitution coefficient $\varepsilon_{\text{s.s.}}$ calculated from Eq. (5) with the same δ and steady-state reduced temperature $u_{\text{s.s.}}$ as in the gas of viscoelastic particles, and with the averaged normal impact velocity, $|\mathbf{c}_{12} \cdot \mathbf{e}| = 4/\sqrt{2\pi}$.

In Fig. 5 we show the dependence of Sonince coefficients on (dimensionless) time for larger dissipation, $\delta = 0.3$. The behavior of the Sonine coefficients is qualitatively the same as for smaller δ. Moreover, the subsequent coefficients a_k decrease by an order of magnitude with increasing k, which again indicates the convergence of the Sonine-polynomial expansion for this dissipation. As the same time a significant discrepancy between the kinetic theory and the MD results is obvious, Fig. 5. In the earlier work [38], two of us have reported that the expansion (5) for ε with $N_k = 20$ does not yield an ac-



FIG. 3: Evolution of a_2 and a_3 for the force-free and heated granular gases with $\delta = 0.15$. The symbols correspond to simulation results, the lines to our theory [numerical solution of Eqs. (20)].



FIG. 4: Evolution of Sonine coefficients a_k $(k = 2, 3, 4, 5)$ for a force-free gas. Here the dissipation parameter $\delta = 0.1$. The symbols correspond to the MD results and the lines to the theory.

ceptable accuracy for the Sonine coefficient a_2 and a_3 for $\delta \geq 0.3$. The discrepancy seen in Fig. 5 between the theoretical curves and the simulation data supports this theoretical conclusion.

B. Uniformly heated gas

We study a uniformly-heated granular gas of viscoelastic particles using the algorithm suggested in Ref. [42] (see also [43, 44]). During event-driven simulations, we heat the system after a time-step dt by adding to the velocity of each i-th particle a random increment, which mimics the heating by noise,

$$v_{i,\alpha}(t + dt) = v_{i,\alpha}(t) + \sqrt{r}\sqrt{dt}\varphi \, , \tag{28}$$

where $\alpha = x, y, z$. In Eq. (28), φ is a random number uniformly distributed within interval $[-0.5, 0.5]$, and r is the amplitude of the noise, $r = 12\xi_0^2$. After the change in velocities, the system is transferred to the centre-of-mass frame,

$$\mathbf{v}_i = \mathbf{v}_i - \frac{1}{N}\sum_{i=1}^{N} \mathbf{v}_i \, ,$$

which ensures conservation of the momentum. In our simulations $dt = 0.1$ and $r = 0.1$ were taken. We used the random kicking interval dt small, as compared to the mean free time. The number of particles and the system size were the same as in the force-free simulations; the initial conditions have been also prepared as in the force-free case. All statistical quantities were obtained as averages over 10 independent runs. The results are shown in Figs. 2, 3, 7 and 6. As it follows from Fig. 2, during the early stages of cooling, the temperature of a heated granular gas evolves in the same way as the respective force-free gas with the same δ. However, while the temperature of the heated gas settles at a steady-state value, the force-free gas continues to cool down. The results of our kinetic theory for $u(\tau)$ perfectly agree with the simulation data. In Fig. 3, we plot a_2, a_3 as functions of the dimensionless time τ for both force-free and heated gas with $\delta = 0.15$. For the initial stage of evolution, at



FIG. 5: Evolution of a_2, a_3, a_4, a_5 for a free granular gas with $\delta = 0.3$.

$\tau \leq 100$, the Sonine coefficients for heated and force-free gases almost coincide. At later times, the coefficients of the heated gas settle at the steady-state [cf. Eqs. (26)-(27)], whereas these for the freely evolving gas continue to decay. Again, as for the force-free gases, our theory is again in excellent agreement with computer simulations.

Figure 6 demonstrates the convergence of the Sonine-polynomial expansion for heated granular gases. As in Fig. 4, we plot here the dependence of a_2, a_3, a_4 and a_5 on τ for $\delta = 0.1$. We see that successive Sonine coefficients decrease by an order of magnitude, which clearly indicates the convergence of the Sonine series. Finally, in Fig. 7, we plot a_2, a_3, a_4 and a_5 for the heated gas with relatively large dissipation, $\delta = 0.3$. In this case one can see a noticeable discrepancy between our theory and the numerical results (cf. Fig. 5 for the freely-evolving gas). Nevertheless, smallness of the high-order coefficients a_4 and a_5 seemingly manifests the convergence of the Sonine series. Interestingly, although the simulation data deviate from the theoretical curves during the relaxation to the steady state, the numerical and theoretical values of the Sonine coefficients a_2 and a_3 in the steady state are very close, see Fig. 7 for $\tau > 10^3$.



FIG. 6: Evolution of Sonine coefficients a_k ($k = 2, 3, 4, 5$) for a uniformly-heated gas. Here, the dissipation parameter $\delta = 0.1$. Gray lines correspond to evolution of granular gas with effective constant restitution coeffcient, calculated according to Eq. (5) with same δ and steady-state relative temperature u, as in the viscoelastic case, and $\langle |\mathbf{c}_{12} \cdot \mathbf{n}| \rangle = 4/\sqrt{2\pi}$.

C. Comparison with the case of a constant restitution coefficient

As we have already mentioned in the Introduction, the behavior of a granular gas with a constant restitution coefficient in a homogeneous cooling state qualitatively differs from that of a gas of viscoelastic particles. The reason for this is the permanent cooling of the gas, which causes the permanent increase of the effective restitution coefficient. Contrary, uniformly heated gases evolve to a steady state, with constant temperature $T_{\rm s.s.}$ where an effective restitution coefficient is kept constant. Therefore, it is natural to assume a somewhat close behavior of the two gases, with the same initial temperature T_0, number density and heating rate, provided the effective restitution coefficient $\varepsilon_{\rm s.s.}$ is appropriately chosen. To chose $\varepsilon_{\rm eff}$ we assume that it should be equal to the restitution coefficient of viscoelastic particles colliding with the relative velocity, equal thermal velocity at the steady state $T_{\rm s.s.}$, with the average normal component of the impact velocity $\langle |\mathbf{c}_{12} \cdot \mathbf{n}| \rangle$. That is, to compute $\varepsilon_{\rm eff}$ we exploit Eq. (5)



FIG. 7: Evolution of a_2, a_3, a_4, a_5 for a uniformly-heated gas with $\delta = 0.3$. Gray lines correspond to evolution of granular gas with effective constant restitution coeffcient, calculated according to Eq. (5) with same δ and steady-state relative temperature u, as in the viscoelastic case, and $\langle |\mathbf{c}_{12} \cdot \mathbf{n}| \rangle = 4/\sqrt{2\pi}$.

with $u(t) = u_{\rm s.s.} = T_{\rm s.s.}/T_0$ and $\langle |\mathbf{c}_{12} \cdot \mathbf{n}| \rangle = 4/\sqrt{2\pi}$, with the corresponding dissipation δ. Fig. 2 demonstrates almost coinciding behavior of the granular temperatures in the two gases. Evolution of temperature towards the steady-state in the in the gas with the constant restitution coefficient $\varepsilon = \varepsilon_{\rm eff}$ is practically indistinguishable from that of the gas of viscoelastic particles; a small discrepancy may be seen only for intermediate times.

Nevertheless, a quantitative agreement in the evolution of the velocity distribution function of the two systems is lacking; only qualitative agreement is observed in the time dependence of the Sonine coefficients. Still however, when the dissipation is not large, $\delta = 0.1$, the steady state values of a_2 and a_3 are almost the same for the two gases, Fig. 6. For larger dissipation, $\delta = 0.3$, not only evolution of these coefficients is rather different, but also their steady-state limits, Fig. 7. Hence we conclude that while for small dissipation it is possible to use an effective constant restitution coefficient to model simultaneously temperature and the form of the velocity distribution function (quantified by a_2 and a_3) in a steady state, this can not be done for large dissipations.

Therefore, as it follows from our study a simplified collision model with an effective constant restitution coefficient may be used only to describe some averaged, that is, macroscopic characteristics, like temperature, but not microscopic, as the velocity distribution function.

IV. CONCLUSION

To date, most studies of granular gases have been focused on systems with a constant restitution coefficient ε, although the basic physics requires impact-velocity dependence of ε. Here we have studied numerically, by means of Molecular Dynamics, and theoretically a granular gas of particles with the impact-velocity dependent restitution coefficient, as it follows from the simplest first-principle model of viscoelastic spheres. To explore evolution of granular temperature as well as the velocity distribution function we performed large-scale MD simulations of a force-free gas and of a gas under a uniform heating. We also developed a theory of the velocity distribution function for a driven gas of viscoelastic particles. The velocity distribution function deviates from a Maxwellian distribution and deviations are quantified by Sonine coefficients in the Sonine polynomials expansion. In our MD simulations we have found a few first Sonine coefficients a_2, a_3, a_4, a_5 and obtained theoretical predictions for a_2 and a_3. We noticed that each next order Sonine coefficient decreases by an order of magnitude with respect to the previous one, that is, a_3 is approximately 10 times smaller than a_2, while a_4 is 10 times smaller than a_3 but 10 time larger than a_5. This implies the convergence of the Sonine series for both heated and force-free gases in the range of dissipation studied.

We observed that predictions of our theory are in a very good agreement with the MD results for granular temperature for both small and large dissipation, quantified by the dissipative parameter δ. We also found that if dissipation is not large, ($\delta = 0.1$), the dependence on time of the Sonine coefficient, obtained in simulations is in excellent agreement with the theoretical predictions for both cases of a homogeneous cooling state (existed theory) and of a uniform heating (the novel theory); at the same time for large dissipation ($\delta = 0.3$) a noticeable deviation between the theory and MD results are observed for.

We also analyze the accuracy of the simplified model of the effective constant restitution coefficient $\varepsilon_{\rm eff}$ for uniformly heated gases. We reveal, that while this model may be used to describe the temperature of a gas, it fails to describe quantitatively both temperature and the velocity distribution function simultaneously, especially for large dissipations.

V. APPENDIX

Here we present numerical values of the coefficients $M_{p,i}^{(k)}$ for p-th moments of the dimensionless collision integral $\mu_p = -\int d\mathbf{c} c^p \tilde{I}(\tilde{f}, \tilde{f})$, where $p = 2, 4, 6$ (Eq. 19). They can be calculated using a formula manipulation program as explained in detail in Ref. [8].

TABLE I: Numerical coefficients for μ_2.

k	$M_{2,0}^{(k)}$	$M_{2,2}^{(k)}$	$M_{2,3}^{(k)}$	$M_{2,22}^{(k)}$	$M_{2,32}^{(k)}$	$M_{2,23}^{(k)}$
0	0	0	0	0	0	0
1	0	0	0	0	0	0
2	6.485666782	1.55656004	0.103770674	0.05447964	0.012347568	0.043583201
3	0	0	0	0	0	0
4	-9.285794108	-2.762523735	-0.138126145	-0.067336539	-0.014198354	-0.051624196
5	-1.801263542	-0.591039599	-0.024626646	-0.011543787	-0.002345029	-0.008657705
6	10.39566154	3.742438128	0.124747864	0.0561366	0.010977733	0.041168185
7	5.911161093	2.32382521	0.058095712	0.025053679	0.004712918	0.017955684
8	-10.43948573	-4.462880197	-0.074381459	-0.030682122	-0.005549036	-0.02147679
9	-10.53072647	-4.877042711	-0.040642187	-0.016002713	-0.002778439	-0.01093409
10	8.770061385	4.385030729	-6.28319E-08	1.5708E-07	-6.9115E-07	-1.09956E-06
11	14.12940508	7.603386089	-0.063361675	-0.022572577	-0.003606668	-0.014669549
12	-4.537693883	-2.620518195	0.043675344	0.014740329	0.00225554	0.009334873
13	-16.3720861	-10.11999574	0.25299992	0.080642707	0.011807124	0.04972868
14	-1.57951827	-1.042482063	0.034749403	0.010424777	0.001458779	0.006254857
15	16.74058856	11.77072632	-0.490447192	-0.137938722	-0.018439173	-0.080466684
16	8.091094766	6.048093381	-0.302404622	-0.079381265	-0.010120669	-0.044983133
17	-14.28935468	-11.33324447	0.661105967	0.161144633	0.019577921	0.088628216
18	-13.91371204	-11.68751806	0.779168081	0.175313108	0.020253112	0.093501317
19	8.817864446	7.831365892	-0.587352513	-0.12114108	-0.013292359	-0.062590324
20	18.12640747	16.99350704	-1.416125536	-0.265523424	-0.027660643	-0.132762689

TABLE II: Numerical coefficients for μ_4.

k	$M_{4,0}^{(k)}$	$M_{4,2}^{(k)}$	$M_{4,3}^{(k)}$	$M_{4,22}^{(k)}$	$M_{4,32}^{(k)}$	$M_{4,23}^{(k)}$
0	0	10.0265131	-2.506628274	0.313328534	0.048957583	0.156664267
1	0	0	0	0	0	0
2	36.31973398	46.85245668	-6.501232013	-0.294188929	-0.014744639	-0.137283095
3	0	0	0	0	0	0
4	-71.50061454	-100.6607515	16.08697411	0.588254117	0.028578585	0.254896238
5	-10.35726536	-15.38813811	2.65528065	0.077425377	0.003372684	0.032260122
6	116.2110953	170.7289911	-29.24466734	-0.978197119	-0.053925433	-0.391290111
7	45.69327521	73.08622368	-13.9966349	-0.27504694	-0.011868725	-0.105434433
8	-169.5076903	-260.3837618	47.12363979	1.010942243	0.055144437	0.370717555
9	-117.6984195	-196.0891402	39.54174543	0.343236574	0.015776747	0.120152601
10	219.5627451	354.2742165	-67.35573679	9.42478E-07	-9.59128E-05	2.19911E-06
11	234.6065359	406.4486348	-85.0337586	0.942565153	0.041890374	0.298516588
12	-240.6229512	-406.3774129	79.91582806	-2.489342054	-0.113668109	-0.746801174
13	-398.7161877	-724.5838699	157.8564548	-6.21062323	-0.248019053	-1.759667007
14	198.2681578	345.5995343	-67.77893928	5.334853916	0.215424796	1.422686368
15	597.9839776	1147.255623	-261.5391558	19.23940936	0.663323094	4.809946224
16	-57.48669811	-83.77829734	7.187455294	-3.892354379	-0.14905002	-0.908182286
17	-797.173725	-1618.265223	386.3071566	-43.41915585	-1.250041435	-9.407465346
18	-208.1541912	-471.7945326	132.0933869	-11.561012	-0.264278169	-2.312272525
19	935.7389044	2010.206189	-501.6884258	77.95690697	1.814066361	14.29254849
20	604.5584047	1387.679757	-379.5240628	55.87757953	1.100691783	9.313324364

TABLE III: Numerical coefficients for μ_6.

k	$M_{6,\,0}^{(k)}$	$M_{6,\,2}^{(k)}$	$M_{6,\,3}^{(k)}$	$M_{6,\,22}^{(k)}$	$M_{6,\,32}^{(k)}$	$M_{6,\,23}^{(k)}$
0	0	112.7982724	-84.5987043	-4.934924416	0.022030913	-0.822487403
1	0	0	0	0	0	0
2	209.9410334	633.7845455	-245.0212299	16.75084546	0.263561766	2.233405168
3	0	0	0	0	0	0
4	-525.042013	-1718.358426	717.5389515	-48.5267813	-0.447724986	-4.852819306
5	-61.2148156	-203.8361017	86.29524105	-7.201298665	-0.061076884	-0.600095271
6	1097.056804	3731.587123	-1607.525698	133.8829794	0.639814941	8.92527426
7	338.2152097	1206.681042	-544.0572385	46.03791709	0.181904962	2.302036496
8	-2042.435348	-7140.741695	3126.21351	-347.473825	-0.750086425	-11.58235962
9	-1116.580388	-4130.985458	1913.431002	-196.4184963	-0.205720952	-3.273691172
10	3411.978481	12219.71999	-5395.763075	806.6588655	-0.002230531	-0.001201659
11	2856.108247	10832.88409	-5076.69307	683.3582205	-0.646810061	-11.39021812
12	-5055.198635	-18485.31829	8165.55863	-1634.92685	3.291775252	54.49778445
13	-6187.93005	-24012.30452	11320.12826	-2006.585914	5.40301205	100.3307334
14	6435.915735	23886.43086	-10416.97776	2818.339574	-10.83459224	-187.8895259
15	11807.48498	46936.94502	-22224.52202	5062.759585	-21.37481894	-421.907027
16	-6433.236775	-23805.35645	9879.364245	-3867.079426	21.1398954	386.7062636
17	-20203.41898	-82418.70275	39185.83747	-11160.63025	60.2704598	1302.074732
18	3203.277562	10083.97492	-2516.563404	3251.164745	-23.50300074	-433.4876022
19	31188.58356	130743.2016	-62400.7815	21721.86788	-133.1393616	-3258.298424
20	5784.30056	29433.94954	-17812.70067	2307.910281	-6.449835295	-384.6515338

---

[1] Physics of Dry Granular Media, edited by H. J. Herrmann, J.-P. Hovi, and S. Luding, NATO ASI Series (Kluwer, Dordrecht, 1998).
[2] H.M. Jaeger, S.R. Nagel, R.P. Behringer, Rev. Mod. Phys. **68**, 1259 (1996).
[3] H. Hinrichsen, D.E. Wolf, *The Physics of Granular Media* (Wiley, Berlin, 2004).
[4] G.H. Ristow, *Pattern Formation in Granular Materials* (Springer-Verlag, Berlin, 2000).
[5] J. Duran, *Sands, Powders and Grains* (Springer-Verlag, Berlin, 2000).
[6] Granular Gases, edited by T. Poeschel and S. Luding, Lecture Notes in Physics Vol. 564 (Springer, Berlin, 2001).
[7] Granular Gas Dynamics, edited by T. Poeschel and N. V. Brilliantov, Lecture Notes in Physics Vol. 624 (Springer, Berlin, 2003).
[8] N.V. Brilliantov, T. Pöschel, *Kinetic Theory of Granular Gases* (Oxford University Press, Oxford, 2004).
[9] I. Goldhirsch, Ann. Rev. Fluid Mech. **35**, 267 (2003).
[10] Planetary Rings, edited by R. Greenberg and A. Brahic (Arizona University Press, Tucson, 1984).
[11] P.K. Haff, J. Fluid Mech. **134**, 401 (1983).
[12] I. Goldhirsch, G. Zanetti, Phys. Rev. Lett. **70**, 1619 (1993).
[13] S.K. Das, S. Puri, Europhys. Lett. **61**, 749 (2003).
[14] S.K. Das, S. Puri, Phys. Rev. E **68**, 011302 (2003).
[15] H. Nakanishi, Phys. Rev. E **67**, 010301 (2003).
[16] T.P.C. van Noije, M.H. Ernst, R. Brito, J.A.G. Orza, Phys. Rev. Lett. **79**, 411 (1997).
[17] T.P.C. van Noije, M.H. Ernst, R. Brito, Phys. Rev. E **57**, R4891 (1998).
[18] S.R. Ahmad, S. Puri, Europhys. Lett. **75**, 56 (2006).
[19] S.R. Ahmad, S. Puri, Phys. Rev. E **75**, 031302 (2007).
[20] J.M. Montanero, A. Santos, Granular Matter **2**, 53 (1998).
[21] T.P.C. van Noije and M.H. Ernst, Granular Matter **1**, 57 (1998).
[22] W. Goldsmit, *The Theory and Physical Behavior of Colliding Solids* (Arnold, London, 1960).
[23] F.C. Bridges, A. Hatzes, D.N.C. Lin, Nature **309**, 333 (1984).
[24] G. Kuwabara, K. Kono, J. Appl. Phys. Part 1 **26**, 1230 (1987).
[25] T. Tanaka, T. Ishida, Y. Tsuji, Trans. Jap. Soc. Mech. Eng. **57**, 456 (1991).
[26] R. Ramirez, T. Pöschel, N.V. Brilliantov, T. Schwager, Phys. Rev. E **60**, 4465 (1999).
[27] N.V. Brilliantov, F. Spahn, J.M. Hertzsch, T. Pöschel, Phys. Rev. E **53**, 5382 (1996).
[28] W.A.M. Morgado, I. Oppenheim, Phys. Rev. E **55**, 1940 (1997).
[29] T. Schwager, T. Pöschel, Phys. Rev. E **57**, 650 (1998).
[30] T. Schwager, T. Pöschel, Phys. Rev. E **78**, 051304 (2008).
[31] A. Goldshtein, M. Shapiro, J. Fliud Mech. **282**, 75 (1995).

[32] S.E. Esipov, T. Pöschel, J. Stat. Phys. **86**, 1385 (1997).
[33] T.P.C. van Noije, M.H. Ernst, Granular Matter **1**, 57 (1998).
[34] N.V. Brilliantov, T. Pöschel, Europhys. Lett. **74**, 424 (2006).
[35] M. Huthmann, J.A.G. Orza, R. Brito, Granular Matter **2**, 189 (2000).
[36] J.J. Brey, M.J. Ruiz-Montero, D. Cubero, Phys. Rev. E **54**, 3664 (1996).
[37] N.V. Brilliantov, T. Pöschel, Phys. Rev. E **61**, 5573 (2000).
[38] A.S. Bodrova, N.V. Brilliantov, Physica A **388**, 3315 (2009).
[39] T. Pöschel, T. Schwager, *Computational Granular Dynamics* (Springer, Berlin, 2005).
[40] N. Brilliantov, C. Saluena, T. Schwager, and T. Poschel, Phys. Rev. Lett. **93**, 134301 (2004).
[41] M.P. Allen, T.J. Tildesley, *Computer Simulation of Liquids* (Oxford University Press, Oxford, 1987).
[42] D.R.M. Williams, F.C. MacKintosh, Phys. Rev. E **54**, R9 (1996); D.R.M. Williams, Physica A **233**, 718 (1996).
[43] T. P. C. van Noije, M. H. Ernst, E. Trizac and I. Pagonabarraga, Phys. Rev. E **59**, 4326 (1999).
[44] J. M. Montanero and A. Santos, Granular Matter, **2**, 53 (2000).